

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/46	A2	(11) International Publication Number: WO 99/04765 (43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/EP98/04517 (22) International Filing Date: 23 July 1998 (23.07.98) (30) Priority Data: MI97A001746 23 July 1997 (23.07.97) IT (71) Applicant (for all designated States except US): CHIESI FARMACEUTICI S.P.A. [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CHIESI, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). VENTURA, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). MEZZADRI, Rosa [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). BRAMBILLA, Gaetano [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). ACERBI, Daniela [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING AN EFFERVESCENT ACID-BASE COUPLE (57) Abstract A pharmaceutical composition in form of effervescent tablets comprising an active ingredient and an effervescent blend, wherein the effervescent blend comprises an acidic component and sodium glycine carbonate as alkaline components. Preferred acid component are fumaric acid, maleic acid and their salts. Tablets are prepared by direct compression in normal thermo-hygrometric conditions and with standard tableting equipment.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PHARMACEUTICAL COMPOSITIONS CONTAINING AN EFFERVESCENT
ACID-BASE COUPLE

This invention relates to orally administered, solid, fast-soluble pharmaceutical compositions containing an effervescent acid-base couple, suitable for dissolving in water or an aqueous solution and for sucking.

The effervescent tablets occupy an important position among dosage forms, being the form of choice not only for adults but also for children. Many drugs, and in particular analgesics, vitamins and antibiotics were designed in this kind of formulations.

The effervescent tablets, when added to cold water, generate a gas which causes effervescence and produces a clear sparkling solution. The gas which gives the effervescence is always carbon dioxide which derives from the reaction between an acid and a base like carbonate or bicarbonate.

The effervescent tablet consists of at least three components:

- the active ingredient;
- an acid;
- an alkali compound (basic ingredient) constituted by a carbonate or a bicarbonate.

The acid and the alkali are the essential components which provide the effervescence and the disintegration of the tablet when is contacted with water.

As acidic component the citric acid both in the hydrated and anhydrous forms is more often used, but other edible acids like tartaric, fumaric, adipic, malic acid can be used as well.

The carbonate, which represents the source of carbon dioxide which generates the effervescence, generally is a water-soluble alkaline carbonate. The choice of the carbonate is very important since, besides provoking the effervescence, it can influence the stability of the tablet.

Sodium bicarbonate is one of the most used carbonate because it is very soluble and of low cost. Alternatively, modified sodium bicarbonate can be used, obtained by heating common sodium bicarbonate in order to convert the surface of its particles to sodium carbonate so increasing its stability.

CONFIRMATION COPY

Other physiologically acceptable alkaline or alkaline earth metal carbonates may be used, such as potassium or calcium (bi)carbonate, sodium carbonate or sodium glycine carbonate.

Compositions of effervescent tablets may also include a lubricant which has to be necessarily selected from the totally water soluble compounds forming a clear solution. Examples of this kind of lubricants are sodium benzoate, sodium acetate, fumaric acid, polyethylenglycols (PEG) higher than 4000, alanine and glycine.

Conventional excipients such as diluents, ligands, bufferings, sweeteners, flavourings, colourings, solubilizers, disintegrants, wetting agents and other excipients of common use may be added to the formulation.

Effervescent tablets are convenient, attractive, easy to use premeasured dosage forms. These advantages, however, are balanced by some technological problems, the two most important of which are hygroscopicity and lubrication.

The instability of effervescent tablets, their tendency to absorb moisture and lose reactivity are generally known. Due to this instability in the presence of water, conventional wet granulation and the subsequent granulate compression are very hardly applicable.

Sometimes the granulation has been carried out using very low amounts of water, for example through the fusion of the citric acid monohydrate, which upon heating releases part of the water of crystallisation which acts as the granulating fluid. Then the granulate has to be processed in conditions of severely controlled relative humidity, normally lower than 20%.

Alternatively, techniques of anhydrous granulation, in the absence of aqueous phases, have been applied using volatile organic solvents like ethanol. However such techniques require special manufacturing environments with strictly controlled relative humidity conditions (normally lower than 20%) and with explosion proof equipment.

Another technique, which is more time-consuming and more laborious, is represented by the separated wet granulation of acidic and alkali granules, which are subsequently mixed and compressed to give the final pharmaceutical composition.

The direct compression of the simple physical blend of the components of the formulation represented an attempt to obviate the above technological difficulties. However such an operation has been carried out in controlled thermo-hygrometric conditions, for example at temperatures lower than 20-25°C and with relative humidity lower than 30%, using tableting machines with tapered dies and punches faced with chromium alloys.

Because of the operating and stability problems, this type of manufacturing method cannot be easily applied to the preparation of effervescent tablets of particular active ingredients which cannot be wet granulated or which contain a residual percentage of water of crystallisation which is hardly eliminated. Typical examples of this kind of medicaments are drug cyclodextrin complexes, hydrated active ingredients and their salts, which may present stability problems in the presence of water. Similar problems are encountered when the composition contains excipients having hydration water or residual-hardly eliminable moisture. Typical examples of this kind of excipients are cyclodextrins .

The other important technological problem affecting the manufacture of effervescent tablets is lubrication, as the lubricant must not only have lipophilic properties for good lubrication, but also high water solubility, to give adequate disintegration and produce quickly a clear solution. Most substances used as lubricants, such as magnesium stearate, are effective, but are water insoluble. The resulting solution after disintegration is cloudy and often has a soapy taste. Ideally, non toxic lubricants with high water solubility and acceptable taste are required. Moreover, the effervescent base is inherently difficult to lubricate, partly due to the nature of the raw materials used and partly due to the rapid tablet disintegration usually required which limits the use of high percentage of lubricants.

Now it has been found, and this is an object of the present invention, that effervescent tablets can be prepared through simple techniques which have direct-industrial application and which are based on the use of a particular effervescent blend of acids and sodium glycine carbonate, provided in a sufficient amount to rapidly disperse and assist dissolution of the components of the formulation.

CONFIRMATION COPY

In particular, according to a second aspect of the invention, it was found that the use of a blend of certain acids with sodium glycine carbonate allows to prepare effervescent tablets by direct compression in normal thermohygrometric conditions and with standard tableting equipment.

It has been even more surprisingly found that this technology applies also to active ingredients and/or excipients which cannot be wet-granulated or which contain a residual percentage of hardly eliminable crystallisation water.

According to a further aspect of the invention, it was found that the use of certain acids/sodium glycine carbonate blend is particularly advantageous in the preparation of effervescent tablets which contain a cyclodextrin as the component of an inclusion complex or as an excipient, despite of the fact that cyclodextrins have hydration water and tend to absorb moisture very easily.

The characteristics of excipients which can be used in the preparation of effervescent tablets, have been described in Aiache JM, Pharm Acta Helv 49 (5/6), 169-178, 1974 and in Boymond C, Labo-Pharma Probl Tech 25(271), 987-995, 1977.

Anyway, Faguet JP et al. in Labo-Pharma Probl Tech 26(274), 207-210, 1978, after evaluating the effects of moisture on the stability of acids, carbonates and bicarbonates, conclude that when sodium glycine carbonate, which is per se moisture - sensitive, is blended with an acid, specifically citric acid, the resulting carbonate is very moisture - unstable, much more than sodium glycine carbonate alone.

In some patents, sodium glycine carbonate is simply mentioned, among various different excipients, as a possible component of effervescent combinations which can be used in chewable tablets (EP 396335), in formulations which form a suspension when contacted with water (EP 528846), in compositions constituted by separated acidic and alkali granules which can also comprise a moisture scavenger (ZA 9307745), in oral, cold-water soluble formulations of β -cyclodextrin complexes with non steroidal anti inflammatory drugs like ibuprofen, naproxen or ketoprofen (WO 9504528).

However, none of the above documents teaches the preparation of effervescent tablets using sodium glycine carbonate as the basic component, nor suggests the possible advantages thereof.

The use of sodium glycine carbonate in effervescent formulations is described in patent applications and scientific literature regarding formulations containing hydrated amoxycillin (PCT WO 9115197), isosorbide-5-mononitrate (DE 4416769) and enzymes (FR 2305194). The acidic component in these formulations is constituted of citric, tartaric, malic or adipic acid and the manufacturing process foresees steps of slugging, milling slugs, blending and compressing or the use of anhydrous excipients or, also, an external lubrication of the machine is performed.

Amela J. et al in the paper "Drug Dev Ind Pharm 22(5), 407-16, 1996" make the analysis of various components which can be used in the preparation of effervescent tablets and they conclude that sodium glycine carbonate is one of the carbonates which do not have favourable compressing characteristics.

The formulations of the invention essentially comprise:

- an active ingredient;
- sodium glycine carbonate;
- an acid capable to react rapidly with sodium glycine carbonate to release carbon dioxide.

One of the preferred acids is fumaric acid which may be present in the form of salt such as mono sodium or potassium fumarate. Certain kinds of formulation take advantage by the lubricant properties of fumaric acid allowing to limit the quantity of lubricant.

Another preferred acid is maleic acid eventually present as a salt.

The choice of the acid is made according to the characteristics of the active ingredient. In some cases mixtures of acids and/or salts are particularly suitable to modulate either the strenght of the acid or the lubricant properties.

The use of fumaric acid in effervescent formulations is described in several documents which refer to various formulations, but never in combination with sodium glycine carbonate.

EP 443381, FR 2715849, WO 9300886, WO 9107174, WO 9104757 are examples of patent literature which mention fumaric acid among other acids which are commonly used in the effervescent pharmaceutical forms such as US 4153678, US 4812303 and US 4704269, referred to formulations of particular active ingredients. In other documents (among which for example GB 1178294, Roscheisen G. and Schmidt PC Eur J. Pharm. Biopharm 41(5), 302-308, 1995), fumaric acid is considered as a lubricant.

Maleic acid has been also described as an acidic component of effervescent couples but never in combination with sodium glycine carbonate.

In a particular embodiment of the invention the active ingredient of the formulation contains residual percentage of moisture or of crystallization water hardly eliminable. Examples of this kind of active ingredients are complexes of drugs such as the piroxicam- β -cyclodextrin complex, levodopa methyl ester and carbidopa hydrate.

The piroxicam- β -cyclodextrin complex has been described in EP 153998, wherein also an effervescent tablet formulation is exemplified. In this case citric acid - sodium glycine carbonate represented the effervescent blend. However, tablets corresponding to the formulation exemplified in EP 153998, have unfavourable characteristics like the opacity of the produced solution, high dimension and weight and low flowability of the pondered blend to be compressed. Moreover, the presence of saccharose as diluent, and of sweetening agents, compromises the stability of the same formulation, as it has been afterwards ascertained.

The formulation may comprise other excipients like:

- a lubricant selected from PEG higher than 4000 and preferably PEG 6000 sodium benzoate, sodium and potassium fumarate, leucine, alanine;
- a sweetening agent selected from aspartame, saccharin, cyclamate, sugars, preferably aspartame;
- a diluent selected from lactose, mannitol, sorbitol or mixtures thereof and preferably spray-dried (SD) lactose and optionally aromatizing agents, ligands, preservatives or others.

As a diluent SD lactose is particularly preferred in that it facilitates the blend flowability so improving compressibility and machinability of the formulation.

The particular effervescent blend of the invention, together with the above mentioned additives, allows highly soluble, stable and small-sized effervescent tablets to be prepared, by direct compressing the component mixture which can be worked at the standard thermo-hygrometric conditions of normal pharmaceutical production facilities, using standard compressing machines with normal punches and dies. Also the subsequent processing, storage and packaging of the tablets can be performed at normal temperature and moisture conditions.

The effervescent compositions of the invention solubilize on contact with water and produce a clear solution for oral administration. Solutions are favoured over suspensions for oral administration, since drugs in solution are more rapidly absorbed. Solutions are also often more acceptable to patients, in terms of palatability. Nevertheless in some cases the active ingredient does not dissolve and the composition does not result in a clear solution, but a suspension. For this kind of active ingredient the possibility to prepare a tablet by direct compression and obtain a rapid disintegration anyway represents a remarkable formulation improvement.

Other advantages of the composition are the low content of sodium ions, due to the employ of sodium glycine carbonate, with respect to other sodium carbonates and the less fizzy effervescence, more pleasant to the patient.

Moreover the composition of the invention, because of its small size, light effervescence and rapid disintegration, can also be prepared as fast dissolving or sucking in the mouth. Infact, as introduced in the mouth, when in contact with saliva, the tablet disintegrates and rapidly forms a solution or an aqueous dispersion easily swallowable.

The following examples further illustrate the invention.

Example 1

Composition of an effervescent tablet having a piroxicam content of 20 mg.

Piroxicam- β -cyclodextrin complex (1:2,5)	191.2 mg
Sodium glycine carbonate	260.0 mg
Fumaric acid	180.0 mg
PEG 6000	20.0 mg
Lactose spray-dried (SD)	208.8 mg
Lemon flavour	25.0 mg
Aspartame	15.0 mg

CONFIRMATION COPY

Piroxicam- β -cyclodextrin, lactose SD, sodium glycine carbonate, lemon flavour, aspartame and PEG 6000 are sieved and pre-mixed. Fumaric acid is added and the components are mixed until an homogeneous blend is obtained. Then the blend is compressed in a standard rotary tableting-machine equipped with round chromium plated punches. The process is carried out at room temperature and with a relative humidity not higher than 55-60%. The dimensions of the produced tablets are about 13 mm diameter, 5 mm thickness and the weight is about 900 mg.

Example 2 - Active ingredient solution test.

The analysis of the tablets prepared according to the example 1 was carried out in order to determine the percentage of the dissolved active ingredient at the end of the effervescence. The maximum dissolution time with effervescence is 1.5 min. The experimental conditions simulated the intake of the effervescent tablets by the patient.

The effervescent tablet was dissolved in three kinds of water. At the end of the effervescence (1.5 min) the amount of piroxicam- β -cyclodextrin was determined in the solution.

The data obtained, which are reported in table 1, demonstrate that the active ingredient concentration in the solution is always higher than 70% of the nominal content per tablet.

Table 1: Dissolution of piroxicam- β -cyclodextrin in 50 ml water

Kind of water	% active ingredient dissolved
demineralized water	72.4
drinking water	85.6
natural no-gassed water	77.0

Example 3 - Dissolution rate of effervescent tablets containing the piroxicam- β -cyclodextrin complex in comparison with standard tablets.

The dissolution rate of effervescent tablets prepared in Example 1 was compared with that of standard piroxicam- β -cyclodextrin tablets using USP Apparatus 2 (paddles) in distilled water at a temperature of 37°C.

	Dissolution time (minutes)	Dissolved piroxicam (%)
20 mg standard tablets	5 10	65% 100%
20 mg effervescent tablets	5	100%

Example 4

The oral absorption profile of piroxicam released from effervescent tablets of piroxicam- β -cyclodextrin complex (β -CD) prepared according to the present invention was compared to that of piroxicam- β CD commercially available standard tablets. The test was carried out after single oral dose administration of the two formulations, equivalent to 20 mg piroxicam, in sixteen healthy volunteers according to a randomized two - way crossover design.

The results, reported in table 2, confirmed the behaviour of the two compositions in the dissolution test showing a more rapid absorption of the active ingredient after administration of the effervescent formulation of the invention. Compared to the standard formulation, the effervescent tablet gives remarkably higher plasma concentrations (Cp of 1.93 μ g/ml vs 0.77 μ g/ml, respectively) 15 min after the administration, as well as a higher drug exposure during the first hour after the administration, as it is shown by the AUC data (AUC = Area Under the Curve, i.e. the area under the plasma concentration vs time curve) collected after 1 hour.

CONFIRMATION COPY

Table 2: Main pharmacokinetic parameters (geometric means \pm standard deviation), statistical comparison and standard 90% confidential intervals (90% CI).

Pharmacokinetic parameters	Piroxicam- β -CD tablets (n = 16)	Piroxicam- β -CD effervescent tablets (n = 16)	90% CI p*
Cp15min ($\mu\text{g/mL}$)	0.77 (0.310-1.90)	1.93 (1.33-2.80)	160%-389% 0.003
Cp30min ($\mu\text{g/mL}$)	2.01 (1.47-2.73)	2.26 (1.64-3.10)	100%-127% 0.106
Cp45min ($\mu\text{g/mL}$)	2.15 (1.76-2.62)	2.22 (1.66-2.97)	94%-114% 0.570
Cp 1h ($\mu\text{g/mL}$)	2.08 (1.75-2.46)	2.09 (1.58-2.77)	93%-109% 0.845
Cmax ($\mu\text{g/mL}$)	2.23 (1.83-2.72)	2.35 (1.74-3.16)	96%-116% 0.330
AUC 1h ($\mu\text{g}\cdot\text{h/mL}$)	1.55 (1.19-2.01)	1.88 (1.40-2.53)	107%-138% 0.018

* Significance level calculated from the analysis of the variance (ANOVA)

Cmax = Maximum plasma Concentration

Example 5

The stability of the effervescent blend and tablets prepared according to the example 1 was tested at 25°C and at different relative humidity conditions.

Both the effervescent blend and tablets stored at a relative humidity of 11%, 33%, 52% and 75% showed an increase in weight at the beginning of the study, and afterwards a slight weight decrease (see tables 3 and 4).

This behaviour is due to two correlated phenomena: the moisture absorption and subsequent loss of carbon dioxide.

The moisture absorption prevails during the first days with respect to the carbon dioxide release.

The loss of carbon dioxide however was so low that the effervescence characteristics of the tablets were not influenced.

Table 3: Blend % weight variation at room temperature (about 25°C) at different relative humidity (R.H.) conditions.

R.H.	Days						
	1	2	3	4	7	9	11
11%	0.06	0.09	0.11	0.09	0.08	0.07	0.07
33%	0.22	0.24	0.23	0.22	0.21	0.20	0.20
52%	0.30	0.30	0.29	0.29	0.27	0.27	0.27
75%	0.42	0.42	0.41	0.40	0.35	0.33	0.34

CONFIRMATION COPY

Table 4: Tablets % weight variation at room temperature (about 25°C) at different relative humidity (R.H.) conditions.

R.H.	Days						
	1	2	3	4	7	9	11
11%	0.05	0.10	0.14	0.10	0.08	0.07	0.07
33%	0.19	0.20	0.20	0.18	0.17	0.16	0.16
52%	0.26	0.25	0.24	0.23	0.21	0.21	0.20
75%	0.17	0.15	0.13	0.12	0.08	0.07	0.07

The effervescent blend and tablets according to the present invention absorb a very low quantity of water (3,5% for the blend and 3,0% for the tablets) even in very unfavourable storage conditions, for example 11 days at 75% relative humidity.

In these particularly unfavourable conditions, the chemical, technological and effervescence characteristics of the tablets are not substantially modified, as it results from tables 5 and 6.

Table 5: Moisture per cent values and blend purity after 11 day storage at different relative humidity conditions

Test	Initial value	11 day values			
		11% R.H.	33% R.H.	52% R.H.	75% R.H.
1) Moisture (%)	2.4	3.6	4.1	4.5	5.9
2) Purity	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005

Table 6: Chemical and technological characteristics of the tablets after 11 day storage at different relative humidity conditions

Test	Initial value	11 day values			
		11% R.H.	33% R.H.	52% R.H.	75% R.H.
Effervescence characteristics (disint. \leq 5 min)	conforms	conforms	conforms	conforms	conforms
1) Moisture (%)	2.5	3.3	3.9	4.4	5.5
2) Purity	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005

- 1) The percentage of moisture absorption was determined by the Karl Fischer method
- 2) The purity was estimated determining the percentage of 2-aminopyridine which is the main degradation product of piroxicam.

Examples from 6 to 14.

With analogous process as described in Example 1 the following tablet formulations have been prepared:

Example 6

Ambroxol hydrochloride

a) Ambroxol hydrochloride	30 mg
Lactose SD	800 mg
Sodium Glycine Carbonate	400 mg
Fumaric Acid	260 mg
PEG 6000	40 mg
Aspartame	30 mg

The mixture is compressed directly into tablets with a diameter of 17 mm and a thickness of 5.5 mm.

b) Ambroxol hydrochloride	60 mg
Lactose SD	600 mg
Sodium Glycine Carbonate	400 mg
Maleic Acid	250 mg
PEG 6000	40 mg
Aspartame	30 mg

The mixture is compressed directly into tablets with a diameter of 17 mm and a thickness of 5.0 mm.

Example 7

Paracetamol

Paracetamol (acetaminophen)	500 mg
Sodium Glycine Carbonate	260 mg
Fumaric Acid	180 mg
PEG 6000	10 mg

The mixture is compressed directly into tablets with a diameter of 13 mm and a thickness of 5.1 mm.

Example 8

Paracetamol / Domperidone Maleate combination

Paracetamol (acetaminophen)	500 mg
Domperidone maleate	10 mg
Sodium Glycine Carbonate	400 mg
Fumaric Acid	300 mg
PEG 6000	10 mg
Aspartame	20 mg

The mixture is compressed directly into tablets with a diameter of 15mm and a thickness of 5.7 mm.

Example 9**Nimesulide**

a) Nimesulide	50 mg
Lactose SD	500 mg
Sodium Glycine Carbonate	800 mg
Fumaric Acid	180 mg
PEG 6000	40 mg

The mixture is compressed directly into tablets with a diameter of 17 mm and a thickness of 5.2 mm.

b) Nimesulide	50 mg
Betacyclodextrin	300 mg
Lactose SD	50 mg
Sodium Glycine Carbonate	300 mg
Fumaric Acid	180 mg
PEG 6000	20 mg

The mixture is compressed directly into tablets with a diameter of 13 mm and a thickness of 4.98 mm.

Example 10**Ibuprofen**

Ibuprofen	200 mg
Lactose SD	610 mg
Sodium Glycine Carbonate	600 mg
Fumaric Acid	360 mg
PEG 6000	30 mg

The mixture is compressed directly into tablets with a diameter of 20 mm and a thickness of 5.2 mm.

CONFIRMATION COPY

Example 11**Morniflumate**

Morniflumate	175 mg
Lactose SD	300 mg
Sodium Glycine Carbonate	650 mg
Fumaric Acid	800 mg
PEG 6000	50 mg

The mixture is compressed directly into tablets with a diameter of 20 mm and a thickness of 5.0 mm.

Example 12**Levodopa methyl ester (LDME)**

LDME	314 mg
Lactose SD	146 mg
Sodium Glycine Carbonate	260 mg
Fumaric Acid	180 mg

The mixture is compressed directly into tablets with a diameter of 13mm and a thickness of 5.0 mm.

Example 13**Carbidopa monohydrate**

Carbidopa monohydrate	27 mg
Lactose SD	433 mg
Sodium Glycine Carbonate	260 mg
Fumaric Acid	180 mg

The mixture is compressed directly into tablets with a diameter of 13 mm and a thickness of 5.0 mm.

Example 14

LDME / Carbidopa monohydrate combination

LDME	314 mg
Carbidopa monohydrate	27 mg
Lactose SD	539 mg
Sodium Glycine Carbonate	520 mg
Fumaric Acid	360 mg
PEG 6000	40 mg

The mixture is compressed directly into tablets with a diameter of 17 mm and a thickness of 5.0 mm.

Examples 15, 16 and 17

Dissolution rates of the formulations of the examples 6, 7 and 14 determined with the USP Apparatus 2 (paddles).

Example 15 : Ambroxol Hydrochloride

Conditions: medium = HCl 0.1 N; volume = 750 ml;

speed = 50 rpm (rounds per minute); temperature = 37 °C

time : 5 min

% of dissolved drug: 98%

Example 16: Paracetamol

Conditions: medium = distilled water; volume = 900 ml; speed = 50 rpm;

temperature = 37 °C

Time : 5 min

% of dissolved drug : 90.9 %

Example 17 : LDME/Carbidopa monohydrate combination

Conditions: medium = HCl 0.1 N; volume = 750 ml; speed = 50 rpm;

temperature = 37 °C

Time : 5 min

% of dissolved Carbidopa : 94%

% of dissolved LDME : 99%

CONFIRMATION COPY

Example 18

The oral absorption profile of levodopa and carbidopa released from effervescent tablets of LDME/carbidopa monohydrate combination prepared according to example 14 was compared to that of levodopa/carbidopa monohydrate commercially available standard tablets (Sinemet®).

The study was carried out after single oral dose administration of the two formulations in six healthy volunteers according to a cross-over design.

The results, reported in tables 7 and 8 show a more rapid absorption and an active ingredients higher exposure during the first hours after administration of the effervescent formulation in comparison to the standard commercial formulation.

In the table Cp = plasma Concentration; Cmax = maximum plasma Concentration; Tmax = Time to maximum concentration; AUC1h, AUC2h, AUCt = Area Under the Curve of plasma concentration vs time after 1 hour, 2 hours and total, respectively.

A hand-drawn straight line on a white background, representing a linear relationship. The line is black and extends from the bottom-left towards the top-right, indicating a positive slope. It is drawn with a simple, slightly irregular stroke, characteristic of a hand-drawn sketch.

Table 7 : Main pharmacokinetic parameters of Levodopa (geometric mean \pm standard deviation) after oral administration of effervescent tablets of LDME/Carbidopa monohydrate combination of example 14 vs standard commercial tablets of Levodopa/Carbidopa monohydrate combination (Sinemet®) in six healthy volunteers

Levodopa pharmacokinetic parameters	Sinemet® tablets	LDME/Carbidopa effervescent tablets
Cp 15 min (ng/mL)	1292 \pm 321	2787 \pm 1338
Cp 30 min (ng/mL)	965 \pm 304	1705 \pm 989
Cp 45 min (ng/mL)	1158 \pm 703	1339 \pm 882
Cp 1h (ng/mL)	999 \pm 541	1023 \pm 691
Cmax (ng/mL)	2218 \pm 1289	3000 \pm 1592
Tmax (h)	0.6 \pm 0.3	0.3 \pm 0.2
AUC 1h (ng·h/mL)	986 \pm 466	1683 \pm 1074
AUC t (ng·h/mL)	5473 \pm 4678	5123 \pm 4485

CONFIRMATION COPY

Table 8 : Main pharmacokinetic parameters of Carbidopa (geometric mean \pm standard deviation) after oral administration of effervescent tablets of LDME/Carbidopa monohydrate combination of example 14 vs standard commercial tablets of Levodopa/Carbidopa monohydrate combination (Sinemet®) in six healthy volunteers.

Carbidopa pharmacokinetic parameters	Sinemet® tablets	LDME/Carbidopa effervescent tablets
Cp 30 min (ng/mL)	52 \pm 31	46 \pm 33
Cp 45 min (ng/mL)	52 \pm 32	63 \pm 46
Cp 1 h (ng/mL)	59 \pm 39	66 \pm 45
Cp 1.5 h (ng/mL)	69 \pm 47	72 \pm 46
Cp 2 h (ng/mL)	49 \pm 32	68 \pm 42
Cmax (ng/mL)	75 \pm 51	88 \pm 65
Tmax (h)	2.6 \pm 1.7	1.5 \pm 0.8
AUC 2h (ng·h/mL)	44 \pm 12	105 \pm 72
AUC t (ng·h/mL)	230 \pm 144	255 \pm 168

CLAIMS

1. Pharmaceutical composition in form of effervescent tablet comprising an active ingredient and an effervescent couple, the couple comprising an acidic component and an alkaline component, characterized in that the alkaline component is sodium glycine carbonate.
2. Pharmaceutical composition according to claim 1 characterized in that the acidic component is fumaric acid or its salts.
3. Pharmaceutical composition according to claim 1 characterized in that the acidic component is maleic acid or its salts.
4. Pharmaceutical composition according to claims 1-3 characterized in that the active ingredient is unstable in the presence of water.
5. Pharmaceutical composition according to claims 1-4 comprising an excipient unstable in the presence of water.
6. Pharmaceutical composition according to claims 1-5 further comprising a cyclodextrin.
7. Pharmaceutical composition according to claims 1-6 characterized in that the active ingredient is a piroxicam- β -cyclodextrin complex.
8. Pharmaceutical composition according to claims 1-7 characterized in that the active ingredient is a combination of levodopa methyl ester and carbidopa monohydrate.
9. Pharmaceutical composition according to claims 1-8 further characterized by at least one additional excipient selected from the group consisting of ligands, lubricants, sweeteners, solubilizers, colourings, flavourings, diluents, disintegrants, wetting and mixtures thereof.
10. Pharmaceutical composition according to claims 1-9 wherein the diluent is lactose spray-dried.
11. Pharmaceutical composition in form of effervescent tablet according to claims 1-10 characterized in that the effervescent couple is activated by water or saliva, said tablet being substantially completely disintegrable both in water and directly in the mouth.
12. Pharmaceutical composition according to claims 1-11 characterized in that the effervescent tablets are prepared by direct compression in normal thermo-hygrometric conditions and with standard compressing equipment.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/46		A3	(11) International Publication Number: WO 99/04765
			(43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/EP98/04517 (22) International Filing Date: 23 July 1998 (23.07.98) (30) Priority Data: MI97A001746 23 July 1997 (23.07.97) IT (71) Applicant (for all designated States except US): CHIESI FARMACEUTICI S.P.A. [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CHIESI, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). VENTURA, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). MEZZADRI, Rosa [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). BRAMBILLA, Gaetano [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). ACERBI, Daniela [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 10 June 1999 (10.06.99)
(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING AN EFFERVESCENT ACID-BASE COUPLE			
(57) Abstract A pharmaceutical composition in form of effervescent tablets comprising an active ingredient and an effervescent blend, wherein the effervescent blend comprises an acidic component and sodium glycine carbonate as alkaline components. Preferred acid component are fumaric acid, maleic acid and their salts. Tablets are prepared by direct compression in normal thermo-hygrometric conditions and with standard tableting equipment.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04517

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K9/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 153 998 A (CHIESI FARMACEUTICI S.P.A.) 11 September 1985 cited in the application see page 17, line 28 - page 18, line 7 ---	1,4-7,9, 11,12
X	WO 96 33694 A (YAMANOUCHI EUROPE B.V.) 31 October 1996 see page 11; example 9 see claims 6,8 --- -/--	1,2

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 March 1999

Date of mailing of the international search report

16/03/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Benz, K

INTERNATIONAL SEARCH REPORT

Int l Application No

PCT/EP 98/04517

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>P.C. SCHMIDT ET AL.: "Brausetabletten-eine fast vergessene Arzneiform" DIE PHARMAZIE, vol. 45, no. 2, February 1990, pages 89-101, XP000133032 BERLIN (DD) see page 90, paragraph 2. - page 92, paragraph 3.1.</p>	1-12
A	<p>EP 0 443 381 A (TAKEDA CHEMICAL INDUSTRIES, LTD.) 28 August 1991 see claim 18</p>	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04517

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 153998 A	11-09-1985	AT 52263 T	15-05-1990
		AU 550007 B	27-02-1986
		AU 3892385 A	29-08-1985
		BE 900836 A	15-02-1985
		CA 1234106 A	15-03-1988
		DK 76085 A,B,	23-08-1985
		IE 58231 B	11-08-1993
		JP 1703351 C	14-10-1992
		JP 3070705 B	08-11-1991
		JP 60208979 A	21-10-1985
		PT 79997 B	13-01-1987
		US 4603123 A	29-07-1986
WO 9633694 A	31-10-1996	FR 2733420 A	31-10-1996
		AU 5693296 A	18-11-1996
		FR 2733424 A	31-10-1996
EP 443381 A	28-08-1991	AT 108649 T	15-08-1994
		CA 2036283 A	15-08-1991
		DE 69102900 D	25-08-1994
		DE 69102900 T	26-01-1995
		JP 4210926 A	03-08-1992
		US 5223246 A	29-06-1993